

**413760. IXORA COCCINEA L. Rubiaceae.**

From Thailand. Seed presented August 1973 by W. E. Manis, Chieng Mai, to Subtropical Horticulture Research Unit, Miami, Florida. Numbered October 29, 1976.

M-22495. 'Thai Dwarf'. Manis residence, Chieng Mai. Planted at Miami, Florida, September 6, 1973. Popular, fine-textured ornamental shrub in Thailand.

**413761 and 413762. TRITICUM AESTIVUM L. Poaceae.****Common wheat.**

From Australia. Seed presented by K. J. Symes, Australian Wheat Collection, Tamworth, New South Wales. Received October 27, 1976.

**413761.** AUS 18547. 'Jabiru'. White, spring type.

**413762.** AUS 18548. 'Warimba'. White, spring type.

**413763 to 413766.**

From United States. Plants and seeds collected October 12, 1976 by A. J. Oakes, Germplasm Resources Laboratory, Agricultural Research Center, Beltsville, Maryland. Numbered November 1, 1976.

**413763 and 413764.** *MISCANTHUS SINENSIS* var. *GRACILLIMUS* Hitch. Poaceae.

**413763.** 3172. Laurel, Maryland. Tufted ornamental, very narrow green leaves. Cultivated. Plant.

**413764.** 3173. Laurel, Maryland. Seed of P.I. 413763. Cultivated. Seed.

**413765 and 413766.** *SCIRPUS CYPERINUS* (L.) Kunth Cyperaceae.

**413765.** 3174. Beltsville Agricultural Research Center-East, Beltsville, Maryland. Tufted perennial. Possible ornamental. Wild. Plant.

**413766.** 3175. Beltsville Agricultural Research Center-East, Beltsville, Maryland. Seed of P.I. 413765. Wild. Seed.

**413767 to 413780.**

From Australia. Seed presented by Research Center, Wagga, Wagga, through Commissioner, Soil Conservation Service, Sydney, New South Wales. Received September 20, 1976.

**413767.** *VICIA BENGHALENSIS* L. Fabaceae.

**Purple vetch.**

'Popany'. Derived from C.P.I. 16632 introduced 1952 from Estacao Agronomica Nacional, Sacavem, Portugal. Selected for earliness of maturity and more adapted to lower rainfall areas than available commercial varieties.

**413768 to 413774.** *VICIA SATIVA* L.

**Common vetch.**

**413768.** 'Blanchefleur'. Derived from C.P.I. 20344 introduced 1955 from Ecole Nationale de Agriculture, Montpellier, France. Flowers white, seed brown to reddish brown, rarely mottled.

**413769.** 4535/1. Introduced 1965 from Malta under C.P.I. 38127. Originally classified as *V. angustifolia*. About 80 days from seedling emergence to flowering. Seed average size, olive tan.

**413770.** 4535/2. Introduced 1965 from Malta under C.P.I. 38127. Originally classified as *V. angustifolia*. About 105 days from seedling emergence to flowering. Seed small, olive or olive brown with fine mottling.

**413771.** 4535/3. Introduced 1965 from Malta under C.P.I. 38127. Originally classified as *V. angustifolia*. About 100 days from seedling emergence to flowering. Seed medium small, olive.